



4985 SW 74th Court, Miami, FL 33155 USA  
 Tel: (1) 786-220-0379 Fax: (1) 786-513-2733



## Chicken Fecal Quantification ID

Detection and quantification of the fecal associated Chicken gene biomarker by real-time quantitative Polymerase Chain Reaction (qPCR) DNA analytical technology

**Submitter:** Company A  
**Date Received:** January 2, 2017  
**Report Generated:** January 8, 2017

*ND: Not Detected*  
*DNQ: Detected Not Quantified*

SM #	Sample ID	Analysis Requested	Marker Quantified (copies/100 ml)	DNA Analytical Results
SM-6A00001	Sample 1	Chicken Bacteroidetes ID	6.74E+03	<b>Detected</b>
SM-6A00002	Sample 2	Chicken Bacteroidetes ID	ND	Not Detected
SM-6A00003	Sample 3	Chicken Bacteroidetes ID	ND	Not Detected
SM-6A00004	Sample 4	Chicken Bacteroidetes ID	1.06E+04	<b>Detected</b>
SM-6A00005	Sample 5	Chicken Bacteroidetes ID	DNQ	<b>Detected</b>
SM-6A00006	Sample 6	Chicken Bacteroidetes ID	ND	Not Detected

Limitation of Damages – Repayment of Service Price

It is agreed that in the event of breach of any warranty or breach of contract, or negligence of Source Molecular Corporation, as well as its agents or representatives, the liability of the company shall be limited to the repayment, to the purchaser (submitter), of the individual analysis price paid by him/her to Source Molecular Corp. The company shall not be liable for any damages, either direct or consequential. Source Molecular Corp. provides analytical services on a PRIME CONTRACT BASIS ONLY. Terms are available upon request. The sample(s) cited in this report may be used for research purposes after an archiving period of 3 months from the date of this report. Research includes, but is not limited to internal validation studies and peer-reviewed research publications. Anonymity of the sample(s), including the exact geographic location will be maintained by assigning an arbitrary internal reference. These anonymous samples will only be grouped by state / province of origin for research purposes. The client must contact Source Molecular in writing within 10 days from the date of this report if he/she does not wish for their submitted sample(s) to be used for any type of future research.

**Revision 1.2**  
**Effective Date 11/2/17**



4985 SW 74th Court, Miami, FL 33155 USA  
 Tel: (1) 786-220-0379 Fax: (1) 786-513-2733



## Preliminary Interpretation of Chicken Fecal “Quantification” ID Results

Detection and quantification of the fecal associated Chicken gene biomarker by real-time quantitative Polymerase Chain Reaction (qPCR) DNA analytical technology

**Submitter:** Company A  
**Date Received:** January 2, 2017  
**Report Generated:** January 8, 2017

Sample ID	INTERPRETATION	
	Concentration of Chicken Fecal Pollution in Sample	Comment
Sample 1	<b>Low Concentration</b>	Low levels of Chicken fecal biomarker(s)
Sample 2	Not Detected	Chicken fecal biomarker not detected
Sample 3	Not Detected	Chicken fecal biomarker not detected
Sample 4	<b>Moderate Concentration</b>	Moderate levels of Chicken fecal biomarker(s)
Sample 5	<b>Low Concentration</b>	Low levels of Chicken fecal biomarker(s)
Sample 6	Not Detected	Chicken fecal biomarker not detected

The opinions/interpretations identified/expressed in this report are outside the scope of this organization's A2LA Accreditation.

## Laboratory Comments

Submitter: Company A

Report Generated: January 8, 2017

### **Non-Detect Results**

In sample(s) classified as non-detect, the host-associated fecal gene biomarker(s) was either not detected in test replicates, one replicate was detected at a cycle threshold greater than 35 and the other was not, or one replicate was detected at a cycle threshold less than 35 and the other was not after repeated analysis.

### **Detected Results**

In sample(s) classified as detected, the host-associated fecal gene biomarker(s) was detected in both test replicates suggesting that the host's fecal contamination is present in the sample(s). Copy number measurements reported are relative, not absolute, quantification.

### **Detected Not Quantified (DNQ) Results**

In sample(s) classified as Detected Not Quantified (DNQ), the host-associated fecal biomarker was detected in both test replicates but in quantities below the limit of quantification. This result indicates that fecal indicators associated with the respective host was present in the sample(s) but in low concentrations.

### **Fecal Reference Samples**

The client is encouraged to submit fecal samples from suspected sources in the surrounding area in order to gain a better understanding of the concentration of the host-associated biomarker with the regional population. A more precise interpretation would be available to the client with the submittal of such baseline samples.

### **Result Interpretations**

Quantitative results are reported along with interpretations. Interpretations are given as "non-detect", "low concentration", "moderate concentration", or "high concentration" based on the concentration of the genetic markers found in the sample(s).

The presence of the biomarker does not signify the presence or absence of that form of fecal pollution conclusively. Only repeated sampling will enable you to draw more definitive conclusions as to the contributor(s) of fecal pollution.

### **Additional Testing**

A portion of all samples has been frozen and will be archived for 3 months. The client is encouraged to perform additional tests on the sample(s) for other hosts suspected of contributing to the fecal contamination. A list of available tests can be found at [sourcemolecular.com/tests](http://sourcemolecular.com/tests)

### **DNA Analytical Method Explanation**

**Water Samples:** Each submitted water sample is filtered through 0.45 micron membrane filter(s). Each filter is placed in a separate, sterile 2ml disposable tube containing a unique mix of beads and lysis buffer. The sample is homogenized for 1min and the DNA extracted using the Generite DNA-EZ ST1 extraction kit (GeneRite, NJ), as per manufacturer's protocol. Deviations to these procedures may occur at the client's request.

**Non-Water Samples:** Each non-water sample submitted by the client is processed as per internal laboratory extraction procedures. An extracted DNA sample is proceed directly to PCR analysis. Details available upon request.

Amplifications to detect the target gene biomarker were run on an Applied Biosystems StepOnePlus real-time thermal cycler (Applied Biosystems, Foster City, CA) in a final reaction volume of 20ul sample extract, forward primer, reverse primer, probe and an optimized buffer. All assays are run in duplicate. Quantification is achieved by extrapolating target gene copy numbers from a standard curve generated from serial dilutions of known gene copy numbers.

For quality control purposes, a positive control and a negative control, were run alongside the sample(s) to ensure a properly functioning reaction and reveal any false negatives or false positives.

## Theory Explanation of Chicken Bacteroidetes “Quantification” ID™

The phylum *Bacteroidetes* is composed of three large groups of bacteria with the best-known category being *Bacteroidaceae*. This family of gram-negative bacteria is found primarily in the intestinal tracts and mucous membranes of warm-blooded animals and is sometimes considered pathogenic.

Comprising *Bacteroidaceae* are the genus *Bacteroides* and *Prevotella*. The latter genus was originally classified within the former (i.e. *Bacteroides*), but since the 1990s it has been classified in a separate genus because of new chemical and biochemical findings. *Bacteroides* and *Prevotella* are gram-negative, anaerobic, rod-shaped bacteria that are inhabitants of the oral, respiratory, intestinal, and urogenital cavities of humans, animals, and insects. They are sometimes pathogenic.

Fecal *Bacteroidetes* are considered for several reasons an interesting alternative to more traditional indicator organisms such as *E. coli* and *Enterococci*.<sup>1</sup> Since they are strict anaerobes, they are indicative of recent fecal contamination when found in water systems. This is a particularly strong reference point when trying to determine recent outbreaks in fecal pollution. They are also more abundant in feces of warm-blooded animals than *E. coli* and *Enterococci*. Furthermore, these latter two organisms are facultative anaerobes and as such they can be problematic for monitoring purposes since it has been shown that they are able to proliferate in soil, sand and sediments.

The Chicken Quantification ID™ service is designed around the principle that fecal *Bacteroidetes* are found in large quantities in feces of warm-blooded animals.<sup>2,3,4,5,6</sup> Furthermore, certain categories of *Bacteroidetes* have been shown to be predominately detected in chickens. Within these *Bacteroidetes*, certain strains of the *Bacteroides* genus have been found in chickens. As such, these bacterial strains can be used as indicators of chicken fecal contamination.

One of the advantages of the Chicken Quantification ID™ service is that the entire water is sampled and filtered for fecal *Bacteroidetes*. As such, this method avoids the random effect of culturing and selecting bacterial isolates off a petri dish. This is a particular advantage for highly contaminated water systems with potential multiple sources of fecal contamination.

Accuracy of the results is possible because the method uses PCR DNA technology. PCR allows quantities of DNA to be amplified into large number of small copies of DNA sequences. This is accomplished with small pieces of DNA called primers that are complementary and specific to the genomes to be detected.

Through a heating process called thermal cycling, the double stranded DNA is denatured and inserted with complementary primers to create exact copies of the DNA fragment desired. This process is repeated rapidly many times ensuring an exponential progression in the number of copied DNA. If the primers are successful in finding a site on the DNA fragment that is specific to the genome to be studied, then billions of copies of the DNA fragment will be available and detected in real-time. The accumulation of DNA product is plotted as an amplification curve. The absence of an amplification curve would indicate that the chicken *Bacteroidetes* gene biomarker is not present.

### References

- <sup>1</sup> Scott, Troy M., Rose, Joan B., Jenkins, Tracie M., Farrah, Samuel R., Lukasik, Jerzy **Microbial Source Tracking: Current Methodology and Future Directions**. Appl. Environ. Microbiol. (2002) 68: 5796-5803.
- <sup>2</sup> Bernhard, A.E., and K.G. Field. **Identification of nonpoint sources of fecal pollution in coastal waters by using host-specific 16S ribosomal DNA genetic markers from fecal anaerobes** Appl. Environ. Microbiol. (2000) 66: 1,587-1,594.
- <sup>3</sup> Bernhard, A.E., and K.G. Field. **A PCR assay to discriminate human and ruminant feces on the basis of host differences in Bacteroides-Prevotella genes encoding 16S rRNA**. Applied and Environmental Microbiology. (2000) 66: 4,571-4,574.
- <sup>4</sup> Fogarty, Lisa R., Voytek, Mary **A Comparison of Bacteroides-Prevotella 16S rRNA Genetic Markers for Fecal Samples from Different Animal Species** Appl. Environ. Microbiol. (2005) 71: 5999-6007.
- <sup>5</sup> Dick, Linda K., Bernhard, Anne E., Brodeur, Timothy J., Santo Domingo, Jorge W., Simpson, Joyce M., Walters, Sarah P., and Field, Katharine G. **Host distributions of uncultivated fecal Bacteroidales bacteria reveal genetic markers for fecal source identification** Appl. Environ. Microbiol. (2005) 71: 3184-3191.
- <sup>6</sup> Weidhaas, Jennifer L., Macbeth, Tamzen W., Olsen, Roger L., and Harwood, Valerie J. **Correlation of Quantitative PCR for a Poultry-Specific Brevibacterium Marker Gene with Bacterial and Chemical Indicators of Water Pollution in a Watershed Impacted by Land Application of Poultry Litter** Appl. Environ. Microbiol. (2011) 77: 2094-2102